



WAS gene

Wiskott-Aldrich syndrome

Normal Function

The *WAS* gene provides instructions for making a protein called WASP. This protein is found in all blood cells. WASP is involved in relaying signals from the surface of blood cells to the actin cytoskeleton, which is a network of fibers that make up the cell's structural framework. WASP signaling activates the cell when it is needed and triggers its movement (motility) and attachment to other cells and tissues (adhesion). In white blood cells, which protect the body from infection, this signaling allows the actin cytoskeleton to establish the interaction between cells and the foreign invaders that they target (immune synapse).

Health Conditions Related to Genetic Changes

severe congenital neutropenia

Wiskott-Aldrich syndrome

More than 350 mutations in the *WAS* gene have been found to cause Wiskott-Aldrich syndrome, a condition characterized by a reduced ability to form blood clots and abnormal immune system function (immune deficiency). Most of these mutations lead to the production of an abnormally short, nonfunctional version of WASP or prevent the production of any WASP. As a result, WASP cannot relay signals, which disrupts the function of the actin cytoskeleton in developing blood cells. White blood cells that lack WASP have a decreased ability to respond to their environment and form immune synapses. As a result, white blood cells are less able to respond to foreign invaders, causing many of the immune problems related to Wiskott-Aldrich syndrome. Similarly, when cells that aid blood clot formation (platelets) lack functional WASP, their development is impaired, leading to a reduction in platelet size and early cell death. The impairments of white blood cells and platelets are largely responsible for the immune deficiency and bleeding problems characteristic of Wiskott-Aldrich syndrome.

X-linked thrombocytopenia

More than 60 mutations in the *WAS* gene have been found to cause X-linked thrombocytopenia, a blood disorder characterized by a decrease in the amount and size of platelets, leading to prolonged bleeding episodes. Immune problems such as an increased susceptibility to infections may also occur. Most of these *WAS* gene mutations change single protein building blocks (amino acids) in WASP. Mutations

typically lead to the production of an altered protein that cannot efficiently relay signals from the cell membrane to the actin cytoskeleton. In people with X-linked thrombocytopenia, these signaling problems primarily affect the development of platelets. In some cases, white blood cells are affected. When WASP function is impaired in white blood cells, these cells are less able to respond to foreign invaders and immune disorders are more likely to occur.

Some *WAS* gene mutations cause X-linked thrombocytopenia in some individuals and a related condition called Wiskott-Aldrich syndrome (described above) in others. These mutations usually prevent the production of any WASP. It is unknown why some people with these mutations have the relatively mild features of X-linked thrombocytopenia and others have the severe symptoms of Wiskott-Aldrich syndrome.


other disorders

Mutations in the *WAS* gene can cause a condition called severe congenital neutropenia, which is characterized by low levels of white blood cells (neutropenia) and decreased survival of hematopoietic stem cells. People with this condition tend to have immune deficiency resulting in recurrent infections. *WAS* gene mutations that cause severe congenital neutropenia lead to a change in a region of WASP known as the Cdc42 binding site. This site must attach (bind) to a protein called Cdc42 in order to turn on (activate) WASP. The mutations that cause severe congenital neutropenia lead to a WASP that is always active (constitutively active), even in the absence of the Cdc42 protein. It is not fully understood how a constitutively active WASP leads to the signs and symptoms of severe congenital neutropenia.

Chromosomal Location

Cytogenetic Location: Xp11.23, which is the short (p) arm of the X chromosome at position 11.23

Molecular Location: base pairs 48,683,753 to 48,691,427 on the X chromosome (Homo sapiens Annotation Release 108, GRCh38.p7) (NCBI)



Credit: Genome Decoration Page/NCBI

Other Names for This Gene

- IMD2
- WASP
- WASP_HUMAN
- Wiskott-Aldrich syndrome (eczema-thrombocytopenia)
- Wiskott-Aldrich syndrome protein

Additional Information & Resources

Educational Resources

- Immunobiology (fifth edition, 2001): Defective T-cell signaling, cytokine production, or cytokine action can cause immunodeficiency.
<https://www.ncbi.nlm.nih.gov/books/NBK27109/#A1510>

GeneReviews

- WAS-Related Disorders
<https://www.ncbi.nlm.nih.gov/books/NBK1178>

Scientific Articles on PubMed

- PubMed
<https://www.ncbi.nlm.nih.gov/pubmed?term=%28%28WAS%5BTIAB%5D%29+OR+%28WASP%5BTI%5D%29%29+AND+%28%28Genes%5BMH%5D%29+OR+%28Genetic+Phenomena%5BMH%5D%29%29+AND+english%5Bla%5D+AND+human%5Bmh%5D+AND+%22last+720+days%22%5Bdp%5D>

OMIM

- NEUTROPENIA, SEVERE CONGENITAL, X-LINKED
<http://omim.org/entry/300299>
- WAS GENE
<http://omim.org/entry/300392>

Research Resources

- Atlas of Genetics and Cytogenetics in Oncology and Haematology
<http://atlasgeneticsoncology.org/Genes/WASID42801chXp11.html>
- ClinVar
<https://www.ncbi.nlm.nih.gov/clinvar?term=WAS%5Bgene%5D>
- HGNC Gene Family: Wiskott-Aldrich Syndrome protein family
<http://www.genenames.org/cgi-bin/genefamilies/set/14>

- HGNC Gene Symbol Report
http://www.genenames.org/cgi-bin/gene_symbol_report?q=data/hgnc_data.php&hgnc_id=12731
- NCBI Gene
<https://www.ncbi.nlm.nih.gov/gene/7454>
- UniProt
<http://www.uniprot.org/uniprot/P42768>

Sources for This Summary

- Ancliff PJ, Blundell MP, Cory GO, Calle Y, Worth A, Kempinski H, Burns S, Jones GE, Sinclair J, Kinnon C, Hann IM, Gale RE, Linch DC, Thrasher AJ. Two novel activating mutations in the Wiskott-Aldrich syndrome protein result in congenital neutropenia. *Blood*. 2006 Oct 1;108(7):2182-9. Epub 2006 Jun 27.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/16804117>
- Blundell MP, Worth A, Bouma G, Thrasher AJ. The Wiskott-Aldrich syndrome: The actin cytoskeleton and immune cell function. *Dis Markers*. 2010;29(3-4):157-75. doi: 10.3233/DMA-2010-0735. Review.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/21178275>
Free article on PubMed Central: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3835520/>
- Devriendt K, Kim AS, Mathijs G, Frints SG, Schwartz M, Van Den Oord JJ, Verhoef GE, Boogaerts MA, Fryns JP, You D, Rosen MK, Vandenberghe P. Constitutively activating mutation in WASP causes X-linked severe congenital neutropenia. *Nat Genet*. 2001 Mar;27(3):313-7.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/11242115>
- Imai K, Morio T, Zhu Y, Jin Y, Itoh S, Kajiwarra M, Yata J, Mizutani S, Ochs HD, Nonoyama S. Clinical course of patients with WASP gene mutations. *Blood*. 2004 Jan 15;103(2):456-64. Epub 2003 Sep 11.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/12969986>
- Lutskiy MI, Rosen FS, Remold-O'Donnell E. Genotype-phenotype linkage in the Wiskott-Aldrich syndrome. *J Immunol*. 2005 Jul 15;175(2):1329-36.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/16002738>
- Notarangelo LD, Ochs HD. Wiskott-Aldrich Syndrome: a model for defective actin reorganization, cell trafficking and synapse formation. *Curr Opin Immunol*. 2003 Oct;15(5):585-91. Review.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/14499269>
- Ochs HD, Notarangelo LD. Structure and function of the Wiskott-Aldrich syndrome protein. *Curr Opin Hematol*. 2005 Jul;12(4):284-91. Review.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/15928485>
- Ochs HD. Mutations of the Wiskott-Aldrich Syndrome Protein affect protein expression and dictate the clinical phenotypes. *Immunol Res*. 2009;44(1-3):84-8. doi: 10.1007/s12026-008-8084-3. Review.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/19082760>
- OMIM: WAS GENE
<http://omim.org/entry/300392>

Reprinted from Genetics Home Reference:
<https://ghr.nlm.nih.gov/gene/WAS>

Reviewed: February 2013
Published: March 21, 2017

Lister Hill National Center for Biomedical Communications
U.S. National Library of Medicine
National Institutes of Health
Department of Health & Human Services